
2015 Recommended Immunizations for Children from Birth Through 6 Years Old



NOTE: If your child misses a shot, about vaccines. if you have questions Talk with your child's doctor doctor for the next shot. you don't need to start over just go back to your child's

shown age range.

FOOTNOTES: * Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

SEE BACK PAGE

COK MORE

NFORMATION ON

ACCINE-PREVENTABLE

PISEASES AND THE

VACCINES THAT

REVENT THEM.

Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and

If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need

For more information, call toll free

http://www.cdc.gov/vaccines 1-800-CDC-INFO (1-800-232-4636)



Control and Prevention Centers for Disease **Health and Human Services** U.S. Department of



of Pediatrics American Academy



DEDICATED TO THE HEALTH OF ALL CHILDREN"